

ON-MASS-SHELL RENORMALIZATION OF THE $\langle \bar{t}t \rangle$
CONTRIBUTION TO THE $\Delta I = 1/2$ $s \leftrightarrow d$ SELF-ENERGY TRANSITION

M.R. AHMADY

International Centre for Theoretical Physics, Miramare P.O. Box 586, 34100 Trieste, Italy

V. ELIAS and N.C.A. HILL

*Department of Applied Mathematics, University of Western Ontario, London, Ontario N6A 5B7,
Canada*

ABSTRACT

In $\langle \bar{t}t \rangle$ -scenarios for dynamical electroweak symmetry breaking, the presence of a very large t-quark condensate necessarily generates an $s \leftrightarrow d$ self-energy transition which will contribute exclusively to $\Delta I = 1/2$ strangeness-changing matrix elements. This contribution is calculated and compared to the purely perturbative contribution, which is well-known to be far too small to have an appreciable effect on $\Delta I = 1/2$ strangeness-changing matrix elements after mass-shell renormalization subtractions are taken into account. The ratio between the t-quark condensate's contribution and the standard-model perturbative contribution to the mass-shell renormalized $s \leftrightarrow d$ self-energy transition is shown to be of order $|m_t \langle \bar{t}t \rangle| / M_W^4$, suggesting that the large t-quark condensate proposed for electroweak symmetry breaking may also be responsible for the $\Delta I = 1/2$ enhancement of strangeness-changing decays.

If electroweak symmetry breaking is driven dynamically by a t-quark condensate $\langle \bar{t}t \rangle$, such a condensate would necessarily have an enormous magnitude. A symmetry-breaking scenario based upon a Nambu-Jona-Lasinio mechanism would suggest a condensate of magnitude $-m_t \Lambda^2$, where Λ , the NJL cutoff, is certainly larger than m_t and may even correspond to an order-Planck-mass unification mass scale.^{1,2} Such a large condensate has already been shown not to alter salient electroweak results, such as the less-than-1% deviation of the electroweak ρ -parameter from unity.¹ Nevertheless, it is of genuine importance to examine how such a very large condensate of *standard-model* fermions, as opposed to technifermions outside the purview of CKM mixing, might enter other electroweak results. For example, the presence of a very large t-quark condensate ($\langle \bar{t}t \rangle$) necessarily generates an $s \leftrightarrow d$ self-energy transition (via $\bar{t}s$ and $\bar{s}t$ CKM mixing angles³) which will contribute exclusively to $\Delta I = 1/2$ strangeness-changing matrix elements.

Of course, there already exists a purely perturbative $s \leftrightarrow d$ self-energy transition within the standard model, but this contribution is known to be two to three orders of magnitude too small to have an appreciable effect on $\Delta I = 1/2$ strangeness-changing matrix elements after mass-shell renormalization subtractions are taken into account.⁴ In unitary gauge, the unrenormalized $s \leftrightarrow d$ self-energy amplitude is of the form

$$\Sigma_{ds}(p) = H(p^2) \not{p} (1 - \gamma_5), \quad (1)$$

where $H(p^2)$ contains both constant and momentum-dependent terms (the former being divergent). However, the on-mass-shell renormalization conditions $\bar{u}_d(p)[\Sigma_{ds}(p)]^{ren} = 0$, $[\Sigma_{ds}(p)]^{ren} u_s(p) = 0$, provide sufficient information to determine algebraically the four allowed counterterms contributing to $[\Sigma_{ds}(p)]^{ren} = \Sigma_{ds}(p) + A \not{p} (1 - \gamma_5) + B \not{p} (1 + \gamma_5) + C + D \gamma_5$.⁵ If we approximate $H(p^2)$ with $H(0) + p^2 H'(0)$, we find that⁴

$$\Sigma_{ds}(p)^{ren} = H'(0) \{ [p^2 - m_s^2 - m_d^2] \not{p} (1 - \gamma_5) + m_s m_d [-\not{p} (1 + \gamma_5) + (m_s + m_d) - \gamma_5 (m_s - m_d)] \}, \quad (2)$$

where the t-quark's Fig. 1 (f1) contribution is given by⁴

$$[H'(0)]_{f1} = (G_F / \sqrt{2}) (V_{td})^* V_{ts} (B_t + 5/12) / 4\pi^2, \quad (3)$$

with B_t between -1.0 and -1.3 for m_t between 1.5 M_W and 2.0 M_W . Using the present empirical range of CKM mixing angles $[0.003 \leq |V_{td}| \leq 0.018, 0.030 \leq |V_{ts}| \leq 0.054]$,⁶ we see that $1 \cdot 10^{-11} GeV^{-2} \leq [H'(0)]_{f1} \leq 2 \cdot 10^{-10} GeV^{-2}$. The renormalized transition amplitude (2) can be incorporated into the $\Delta I = 1/2$ component of the $K_s \rightarrow 2\pi$ amplitude,

denoted by ^{4,7}

$$a_{1/2} = 3\sqrt{2}g_{Kqq}(m_s - m_d)M^4(1 - m_\pi^2/m_K^2)H'(0)/(64\pi^2 f_\pi^2) \quad (4)$$

where M is a phenomenological cut-off of order 2 GeV and $g_{Kqq} \approx 3.8$. Using (3) and (4) we find the t-quark's purely perturbative contribution to $a_{1/2}$ to be between $2 \cdot 10^{-9}$ and $9 \cdot 10^{-11}$ GeV, a number at least two orders of magnitude smaller than the experimental value ($3.5 \cdot 10^{-7}$ GeV) reflecting the enhancement of the $\Delta I = 1/2$ component of strangeness-changing nonleptonic decays. A more careful treatment⁸ of the hadronization of $\Sigma_{ds}(p)^{ren}$, based upon a direct comparison of the weak kaon axial-vector current $-i \int d^4x e^{iq \cdot x} \langle 0 | T[A_\mu^3(x) H_W^{\Delta S=1}] | \bar{K}^0 \rangle$ to the strong-interaction axial-vector current $i\sqrt{2}f_{Kq\mu} = -i \int d^4x e^{iq \cdot x} \langle 0 | A_\mu^{6+i7}(x) | \bar{K}^0 \rangle$, leads to virtually the same conclusions: $[H'(0)]_{f1}$ is between two and four orders of magnitude too small to account for the measured $K \rightarrow 2\pi$ decay rate.

If electroweak symmetry breaking is caused by condensation of a t- quark- antiquark pair, residual contributions to the Wick-Dyson expansion of Feynman amplitudes involving the vacuum expectation values of normal ordered t-quark and antiquark fields are no longer seen to vanish. The presence of such contributions is entirely analogous to how $\langle u\bar{u} \rangle$ and $\langle d\bar{d} \rangle$ condensates characterizing the chiral- symmetry breakdown of QCD enter Feynman amplitudes that are utilized in the construction of QCD sum-rules.^{8,9} In particular, the unitary gauge contribution of Fig. 2, corresponding to replacing the time ordered vacuum expectation value of interaction-picture t-quark fields $\langle 0 | T[t(x)\bar{t}(0)] | 0 \rangle$ with the corresponding normal-ordered contribution $\langle 0 | : t(x)\bar{t}(0) : | 0 \rangle$ in the Wick-Dyson expansion of $\langle 0 | T[t_H(x)\bar{t}_H(0)] | 0 \rangle$ [the fermion two-point function of Heisenberg fields], is seen to be

$$\begin{aligned} \Sigma_{ds}(p) &= \frac{G_F M_W^2}{\sqrt{2}} V_{td}^* V_{ts} \int \frac{d^4q}{(2\pi)^4} \int d^4x e^{i(p-q) \cdot x} \\ &\times \left[\frac{g^{\mu\nu} - q^\mu q^\nu / M_W^2}{q^2 - M_W^2} \right] \gamma_\mu (1 - \gamma_5) \langle 0 | : t(x)\bar{t}(0) : | 0 \rangle \gamma_\nu (1 - \gamma_5). \end{aligned} \quad (5)$$

The $\langle \bar{t}t \rangle$ projection of $\langle 0 | : t(x)\bar{t}(0) : | 0 \rangle$ is given by¹⁰

$$\langle 0 | : t(x)\bar{t}(0) : | 0 \rangle = -\frac{\langle \bar{t}t \rangle}{6m_t^2} (i\gamma \cdot \partial + m_t) [J_1(m_t \sqrt{x^2}) / \sqrt{x^2}]. \quad (6)$$

The theoretical consistency of eq. (6) has been examined in a number of places. Insertion of this vacuum expectation value into flavour- diagonal fermion two-point functions has been shown to uphold the gauge invariance of the fermion propagator pole in both covariant¹¹ and noncovariant¹² gauges, despite strong overall

gauge dependence of the fermion self-energy. Moreover, inclusion of the $\langle \bar{f}f \rangle$ projection of the $\langle 0| : f(x)\bar{f}(y) : |0\rangle$ contribution to the Wick-Dyson expansion of the vacuum polarization of a vector-coupled spin-1 field has been shown to yield only transverse contributions, consistent with gauge invariance and the vector-current Ward identity.¹³ Similar inclusion of the $\langle \bar{f}f \rangle$ projection of the $\langle 0| : f(x)\bar{f}(y) : |0\rangle$ -contributions to the Wick-Dyson expansion of the VVA-triangle amplitude has also been demonstrated to uphold the anomalous axial-vector Ward identity.¹⁴

Equation (6) may be re-expressed as follows:¹⁵

$$\langle 0| : t(x)\bar{t}(0) : |0\rangle \equiv \int d^4k e^{-ik \cdot x} (\not{k} + m_t) F_t(k), \quad (7)$$

$$\int d^4k e^{-ik \cdot x} F_t(k) = -\langle \bar{t}t \rangle J_1(m_t \sqrt{x^2}) / (6m_t^2 \sqrt{x^2}), \quad (8)$$

$$\int d^4k e^{-ik \cdot x} g(k^2) F_t(k) = \int d^4k e^{-ik \cdot x} g(m_t^2) F_t(k), \quad (9)$$

where the last property [eq. (9)] is a consequence of $\langle 0| : t(x)\bar{t}(0) : |0\rangle$ being a solution of the free-field Dirac equation. One finds from Eq.(5) that the coefficient $H(p^2)$ in the unrenormalized Fig. 2 (f2) self-energy (1) is given by

$$[H(p^2)]_{f2} = \frac{G_F}{\sqrt{2}} M_W^2 V_{td}^* V_{ts} \left\{ \left(-4 - \frac{2(m_t^2 + p^2)}{M_W^2} \right) \left[\frac{\langle \bar{t}t \rangle}{24m_t p^2} + \frac{p^2 + m_t^2 - M_W^2}{2p^2} I(p^2) \right] + \frac{4m_t^2}{M_W^2} I(p^2) \right\}, \quad (10)$$

$$I(p^2) = \int d^4k F_t(k) / [(p-k)^2 - M_W^2 + i\epsilon]. \quad (11)$$

The integral in Eq. (11) may be evaluated first by exponentiating the propagator and through utilization of Eqs. (9) and (8):¹⁵

$$\begin{aligned} I(p^2) &= -i \int_0^\infty d\eta e^{i\eta(p^2 + m_t^2 - M_W^2 + i\epsilon)} \int d^4k e^{-i(2\eta p) \cdot k} F_t(k) \\ &= \frac{i\langle \bar{t}t \rangle}{6m_t^2} \int_0^\infty d\eta \exp[i\eta(p^2 + m_t^2 - M_W^2 + i\epsilon)] \frac{J_1(2m_t \eta \sqrt{p^2})}{2\eta \sqrt{p^2}} \\ &= -\frac{\langle \bar{t}t \rangle}{24m_t^3 p^2} \left[(p^2 + m_t^2 - M_W^2) - \sqrt{(p^2 - m_t^2)^2 - M_W^2(2p^2 + 2m_t^2 - M_W^2)} \right], \end{aligned} \quad (12)$$

where the final line of Eq.(12) is obtained by use of a tabulated integral.¹⁶ Substitution of the final line of (12) into (10) reveals that the fig. 2 contribution to $H(p^2)$ is analytic at $p^2 = 0$:

$$[H(p^2)]_{f2} = [H(0)]_{f2} + p^2 [H'(0)]_{f2}, \quad (13)$$

$$[H(0)]_{f2} = -\frac{G_F V_{td}^* V_{ts} \langle \bar{t}t \rangle}{4\sqrt{2}(m_t^2 - M_W^2)^2} [1 - 7M_W^2/3m_t^2 - 2M_W^4/3m_t^4], \quad (14)$$

$$[H'(0)]_{f2} = \frac{G_F V_{td}^* V_{ts} \langle \bar{t}t \rangle m_t^5}{12\sqrt{2}(m_t^2 - M_W^2)^4} [1 - 4M_W^2/m_t^2 + 9M_W^4/m_t^4]. \quad (15)$$

The relative contributions of Fig. 2 and Fig. 1 to the on-shell- renormalized self-energy transition, as defined by (2), may be obtained via direct comparison of Eqs. (3) and (15). One finds for m_t between 120 GeV and 160 GeV that $|12m_t \langle \bar{t}t \rangle / M_W^4| \geq |[H'(0)]_{f2} / [H'(0)]_{f1}| \geq |0.4m_t \langle \bar{t}t \rangle / M_W^4|$. As remarked earlier, the Fig. 1 contribution $[H'(0)]_{f1}$ to the $\Delta I = 1/2$ component of the $K \rightarrow 2\pi$ decay amplitude is between two and four orders of magnitude below the experimental value. Consequently, we see that the Fig. 2 contribution $[H'(0)]_{f2}$ yields a contribution *comparable to the experimental amplitude* provided $\langle \bar{t}t \rangle \sim m_t^3$ for maximal mixing angles and $m_t \approx 120 \text{ GeV}$. For minimal mixing angles and $m_t \approx 160 \text{ GeV}$, the value $\langle \bar{t}t \rangle \sim (10m_t)^3$ is compatible with the experimental $K \rightarrow 2\pi$ decay rate.

These results suggest, within the context of t-quark-condensate driven electroweak symmetry breaking, that the large magnitude of the t-quark condensate may be responsible for the $\Delta I = 1/2$ rule. However, these results appear to rule out any sensitivity of the t-quark condensate to Planck-mass unification scales.

References

1. V.A. Miransky, M. Tanabishi and K. Yamawaki, *Phys. Lett.* **B221** (1989) 177.
2. R.R. Mendel, Massachusetts Institute of Technology Ph.D. Thesis, 1982 (unpublished); W.A. Bardeen, C.T. Hill and M. Lindner, *Phys. Rev.* **D41** (1990) 1647.
3. M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49** (1973) 652.
4. S.-P. Chia *Phys. Lett.* **B147**(1984) 361.
5. N.G. Deshpande and G. Eilam *Phys. Rev.* **D26** (1982) 2463.
6. K. Hikasa et. al. (Particle Data Group), *Phys. Rev.* **D45** (1992) S1.
7. B.H.J. McKellar and M.D. Scadron, *Phys. Rev.* **D27** (1983) 157.
8. V. Elias, R.R. Mendel and M.D. Scadron, *Phys. Rev.* **D42** (1990) 1591.
9. M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Nucl. Phys.* **B147** (1979) 385 and 448; P. Pascual and R. Tarrach, *Renormalization for the Practitioner* (Springer-Verlag, Berlin, 1984), pp. 155-191.
10. V. Elias, T.G. Steele and M.D. Scadron, *Phys. Rev.* **D38** (1989) 1584; E. Bagan, J.I. Latorre and P. Pascual, *Z. Phys.* **C32** (1986) 43; F.J. Yndurain, *Z. Phys.* **C42** (1989) 653.
11. V. Elias and M.D. Scadron *Phys. Rev.* **D30** (1984) 647; L.J. Reinders and K. Stam, *Phys. Lett.* **B180** (1986) 125; M.R. Ahmady, V. Elias, R.R. Mendel, M.D. Scadron and T.G. Steele, *Phys. Rev.* **D39** (1989) 2764.
12. V. Elias and T.G. Steele *Phys. Rev.* **D40** (1989) 2669; M. Lavelle and M. Schaden, in *Physical and Nonstandard Gauges*, P. Gaigg, W. Kummer and M. Schweda, eds. (Springer-Verlag, Berlin, 1990) p. 240.
13. T.G. Steele, *Z. Phys.* **C42** (1989) 499.
14. M.R. Ahmady, A.S. Deakin, Mong Tong and V. Elias *Z. Phys.* **C56** (1992) 395.
15. E. Bagan, M.R. Ahmady, V. Elias and T.G. Steele, submitted to *Phys. Rev.* **D**.
16. I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products* (Academic Press, Orlando, 1980) 712 [Eq. 6.623.3].

Figure Captions

Figure 1: Purely perturbative t-quark contribution to the s-d transition amplitude.

Figure 2: $\langle \bar{t}t \rangle$ contribution to the s-d transition amplitude.